

The asymptotic conformal invariance in Chern-Simons theory with matter in curved space-time

Alexandr V. Timoshkin

Tomsk Pedagogical University, 634041 Tomsk, Russia

We consider the asymptotic behaviour of the Chern - Simons (CS) theory with matter in curved space - time. The asymptotics of effective couplings are discussed.

1. The study of renormalizable field theories in curved space [1,2,3,4] proved the existence of the phenomenon of asymptotic conformal invariance. In this short review we discuss the asymptotic behaviour of 3d CS theories in curved space. Let us consider renormalizable abelian CS theory with scalar and spinor in three dimensions [5]. The Lagrangian looks like:

$$L = \frac{1}{2} \epsilon^{\mu\nu\lambda} A_\mu \partial_\nu A_\lambda + |D_\mu \Phi|^2 + i \bar{\Psi} \hat{D} \Psi + \alpha \bar{\Psi} \Psi \Phi^* \Phi - h (\Phi^* \Phi)^3 \quad (1)$$

Here $D_\mu = \partial_\mu - ieA_\mu$, Φ, Ψ -complex scalar and dirac spinor consequently, coupling constants e, h, α are dimensionless. The theory with Lagrangian (1) is multiplicatively renormalizable.

The two - loop RG equation for coupling constants has the form [6,7]:

$$(8\pi)^2 \frac{de(t)}{dt} = 0$$

$$(8\pi)^2 \frac{d\alpha(t)}{dt} = \frac{14}{3} \alpha^3(t) - 34\alpha(t) e^4(t) - 24e^6(t) \quad (2)$$

$$(8\pi)^2 \frac{dh(t)}{dt} = 168h^2(t) - 84h(t) \alpha^2(t) + 36e^8(t) + 8\alpha(t) e^6(t) + 4\alpha^2(t) e^6(t) - 4\alpha^4(t).$$

It has been shown in paper [6] that for the theory with Lagrangian (1) exists finite four cases in which the theory is finite at two - loop level:

$$\begin{aligned} 1. \quad & \alpha = 3e^2, \quad h = -\frac{19}{4}e^4 \\ 2. \quad & \alpha = 3e^2, \quad h = e^4; \end{aligned} \quad (3)$$

$$3. \quad \alpha \approx -2, 23e^2, \quad h \approx -0, 59e^4;$$

$$4. \quad \alpha \approx -2, 23e^2, \quad h \approx 0, 62e^4.$$

In regime of finiteness the effective coupling constants are:

$$e(t) = e, \alpha(t) = \alpha, h(t) = h, \quad (4)$$

where values α, h give one from four variants (3). Let us consider the limit $t \rightarrow \infty$ (infrared) for solutions of equations (2). In this limit independently of initial values

$$h(t) \longrightarrow e^4, \alpha(t) \longrightarrow 3e^2. \quad (5)$$

Therefore the theory becomes finite and supersymmetric in the asymptotic [8]. Consequently supersymmetry is infrared stable as far as in four dimensions [8]. Let us study now the theory in the limit $t \longrightarrow +\infty$. In this case we fix the initial value for α : $\alpha(0) = 3e^2$. Then $\alpha(t) = 3e^2$ and at $t \longrightarrow +\infty$, $h(t) \longrightarrow -\frac{19}{4}e^4$ independent of initial value $h(0)$. Consequently in ultraviolet asymptotic under fixed value $\alpha(0) = 3e^2$ the theory effectively becomes finite (asymptotic finiteness [8]). In this case initial value $h(t)$ is arbitrary. Fix now $\alpha(0) \approx -2, 23e^2$. Then at $t \longrightarrow \infty$, $h(t) \longrightarrow 0, 62e^4$ independently of initial value $h(0)$. The theory again appears to be asymptotic finite. In ultraviolet limit two regimes of asymptotic finiteness exist.

2. Let us consider the behavior of effective charge $\xi(t)$ in abelian CS theory with matter in curved space - time, the beta-function on two - loop level has the form [10].

$$\beta_\xi^{(2)} = \left(\xi - \frac{1}{8} \right) \gamma_{m^2}^{(2)}, \quad (6)$$

where $\gamma_{m^2}^{(2)}$ -gamma-function for mass of the scalar field in two - loop approximation. Then with supposition (6), two - loop RG equation for $\xi(t)$ has the form:

$$\frac{d\xi(t)}{dt} = \left(\xi(t) - \frac{1}{8} \right) \cdot \gamma_{m^2}^{(2)}(t), \quad (7)$$

where $\gamma_{m^2}^{(2)}$ is given in paper [7]. Then the solution of equation (7) in regimes of finiteness (3) has the form:

$$\xi(t) = \frac{1}{8} + \left(\xi - \frac{1}{8} \right) \cdot e^{\gamma_{m^2}^{(2)} \cdot t} \quad (8)$$

where $\gamma_{m^2}^{(2)} = 0$ (for $N = 2$ supersymmetric theory) or $\gamma_{m^2}^{(2)} \approx -\frac{e^4}{24\pi^2} \cdot 11, 23 \times 6, 77$.

Therefore at $\alpha \approx -2, 23e^2$ the theory appears to be asymptotic conformally invariant in $d = 3$ at $t \longrightarrow \infty$, $\xi(t) \longrightarrow \frac{1}{8}$ independent of initial value. At $t \longrightarrow -\infty$, $|\xi(t)| \longrightarrow \infty$. For $N = 2$ there are 18 regimes of finiteness for $\gamma_{m^2}^{(2)}$ can be positive, negative or zero. Then at $t \longrightarrow \infty$ the following situations are possible:

1. $\xi(t) \longrightarrow \frac{1}{8}$ (asymptotic supersymmetry);
2. $\xi(t) = \xi$ (the asymptotic supersymmetry);
3. $|\xi(t)| \longrightarrow \infty$.

For $t \longrightarrow \infty$ the behavior of $\xi(t)$ in 1 and 3 is exchanged. Thus, it is shown that in $d3$ gauge theories also can be asymptotic conformal invariante of exponential type, like in [9]. This short review is mainly based on ref.[10].

Acknowledgements.

A.V.T thanks S.D. Odintsov for help at preparation this article.

References.

- [1]. Buchbinder I.L., Odintsov S.D., Shapiro I.L.//Rivista Nuovo Cim.-1989.-v.12.-p.1; Odintsov S.D.//Fortschr. Phys.- 1991.- v.39.-p.621.
- [2]. Buchbinder I.L., Odintsov S.D.//Sov. Phys. J.-1985.-2116.-p.86; Yad.Phys.-1984.-v.40. -p.1338; Lett. Nuovo Cim.-1985.-v.42.-p.379.
- [3]. Lichtzier I.M., Odintsov S.D.//Yad.Phys.-1986.-v.47.-p.1782; Europhys. Lett.-1988.- v.7.-p.95; Buchbinder I.L., Odintsov S.D., Fonarev O.A.//Sov. Phys.J.-1990.-2116.-p.74; Mod. Phys. Lett.-1989.-v.A4.-p.2713.
- [4]. Buchbinder I.L.//Theor.Math.Phys.-1984.-v.61.-p.393; Fortschr. Phys.-1986.-v.1986.- v.34.-p.605.
- [5]. Deser S., Jackiw R., Templeton S.//Ann. Phys.-1982.-v.140.-p.372.
- [6]. Grigoryev G.V., Kazakov D.I.//Phys. Lett.-1991.-v.253B.-p.372.
- [7]. Avdeev L.V., Grigoryev G.V., Kazakov D.I.//Preprint CERN.-T.H.6091/91,1991.
- [8]. Odintsov S.D., Shapiro I.L.//Lett. JETP.-1989.-v.49.-p.125; Mod.PhysLett.-1989.-v.A4.-p.1479; Odintsov S.D., Shapiro I.L., Toms D.J.//Int. J. Mod Phys.-1991.-v.A6.-p.1829.
- [9]. Buchbinder I.L., Odintsov S.D., Lichtzier I.L.//Theor. Math. Phys.-1989.-v.79.-p.314; Class. Grav.-1989.-v.6.-p.605; Zaripov R.Sh., Odintsov S.D.//Theor. Math. Phys.-1990.- v.83.-p.399; Mod. Phys. Lett.-1989.-v.A4.-p.1955.
- [10] Odintsov S.D., Timoshkin A.V.//Sov. Phys. J.-1992.-2116.-P.45.